

it is represented to contain. The moisture content is not more than 7.5 percent. The methacycline hydrochloride used conforms to the standards prescribed by § 446.50(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The methacycline hydrochloride used in making the batch for potency, moisture, pH, absorptivity, identity, and crystallinity.

(b) The batch for potency and moisture.

(ii) Samples required:

(a) The methacycline hydrochloride used in making the batch: 10 packages, each containing approximately 300 milligrams.

(b) The batch: A minimum of 30 capsules.

(b) *Tests and methods of assay*—(1) *Potency.* Proceed as directed in § 436.106 of this chapter, preparing the sample for assay as follows: Blend a representative number of capsules in a high-speed glass blender jar containing sufficient sterile distilled water to give a stock solution of convenient concentration. Further dilute an aliquot of the stock solution with sterile distilled water to the reference concentration of 0.06 microgram of methacycline per milliliter (estimated).

(2) *Moisture.* Proceed as directed in § 436.201 of this chapter.

[39 FR 19076, May 30, 1974, as amended at 43 FR 11163, Mar. 17, 1978; 46 FR 46313, Sept. 18, 1981; 50 FR 19920, May 13, 1985]

§ 446.150b Methacycline hydrochloride oral suspension.

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity.* Methacycline hydrochloride oral suspension contains methacycline hydrochloride and one or more suitable and harmless buffers, dispersants, diluents, colorings, flavorings, and preservatives. It contains methacycline hydrochloride equivalent to 14 milligrams of methacycline per milliliter. Its potency is satisfactory if it is not less than 90 percent and not more than

125 percent of the number of milligrams of methacycline that it is represented to contain. Its pH is not less than 6.5 nor more than 8.0. The methacycline hydrochloride used conforms to the standards prescribed by § 446.50(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The methacycline hydrochloride used in making the batch for potency, moisture, pH, absorptivity, identity, and crystallinity.

(b) The batch for potency and pH.

(ii) Samples required.

(a) The methacycline hydrochloride used in making the batch: 10 packages, each containing approximately 300 milligrams.

(b) The batch: A minimum of 5 immediate containers.

(b) *Tests and methods of assay*—(1) *Potency.* Proceed as directed in § 436.106 of this chapter, preparing the sample for assay as follows: Transfer an accurately measured representative portion of the well-shaken suspension to an appropriate-sized volumetric flask, and dilute to volume with sterile distilled water. Mix well. Remove an aliquot of the stock solution and further dilute with sterile distilled water to the reference concentration of 0.06 microgram of methacycline per milliliter (estimated).

(2) *pH.* Proceed as directed in § 436.202 of this chapter using the undiluted sample.

[39 FR 19076, May 30, 1974, as amended at 43 FR 11163, Mar. 17, 1978; 50 FR 19920, May 13, 1985]

§ 446.160 Minocycline hydrochloride oral dosage forms.

§ 446.160a Minocycline hydrochloride tablets.

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity.* Minocycline hydrochloride tablets are composed of minocycline hydrochloride and one or more suitable